

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY) 2. REPORT TYPE Technical Papers 3. DATES COVERED (From - To)

4. TITLE AND SUBTITLE 5a. CONTRACT NUMBER

5b. GRANT NUMBER

5c. PROGRAM ELEMENT NUMBER

6. AUTHOR(S) 5d. PROJECT NUMBER

2303

5e. TASK NUMBER

m208

5f. WORK UNIT NUMBER

345709

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 8. PERFORMING ORGANIZATION REPORT

Air Force Research Laboratory (AFMC)

AFRL/PRS

5 Pollux Drive

Edwards AFB CA 93524-7048

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 10. SPONSOR/MONITOR'S ACRONYM(S)

Air Force Research Laboratory (AFMC)

AFRL/PRS

5 Pollux Drive

Edwards AFB CA 93524-7048

11. SPONSOR/MONITOR'S NUMBER(S)

please see attached

12. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for public release; distribution unlimited.

13. SUPPLEMENTARY NOTES

14. ABSTRACT

20030129 229

15. SUBJECT TERMS

16. SECURITY CLASSIFICATION OF:

a. REPORT b. ABSTRACT c. THIS PAGE

Unclassified Unclassified Unclassified

17. LIMITATION OF ABSTRACT

A

18. NUMBER OF PAGES

19a. NAME OF RESPONSIBLE PERSON

Leilani Richardson

19b. TELEPHONE NUMBER (include area code)

(661) 275-5015

CP

MEMORANDUM FOR PRS (In-House Publication)

FROM: PROI (STINFO)

10 April 2001

SUBJECT: Authorization for Release of Technical Information, Control Number: **AFRL-PR-ED-VG-2001-078**
Fajardo, Mario, "Chemistry and Spectroscopy in Solid Parahydrogen"

U. Wyoming Chemistry Dept. Seminar
(Caramie, WY, 20 April 2001) (Deadline: 20 April 2001)

(Statement A)

1. This request has been reviewed by the Foreign Disclosure Office for: a.) appropriateness of distribution statement, b.) military/national critical technology, c.) export controls or distribution restrictions, d.) appropriateness for release to a foreign nation, and e.) technical sensitivity and/or economic sensitivity.

Comments: _____

Signature _____ Date _____

2. This request has been reviewed by the Public Affairs Office for: a.) appropriateness for public release and/or b) possible higher headquarters review

Comments: _____

Signature _____ Date _____

3. This request has been reviewed by the STINFO for: a.) changes if approved as amended, b.) appropriateness of distribution statement, c.) military/national critical technology, d.) economic sensitivity, e.) parallel review completed if required, and f.) format and completion of meeting clearance form if required

Comments: _____

Signature _____ Date _____

4. This request has been reviewed by PRS for: a.) technical accuracy, b.) appropriateness for audience, c.) appropriateness of distribution statement, d.) technical sensitivity and economic sensitivity, e.) military/national critical technology, and f.) data rights and patentability

Comments: _____

APPROVED/APPROVED AS AMENDED/DISAPPROVED

PHILIP A. KESSEL Date
Technical Advisor
Space & Missile Propulsion Division

Chemistry and Spectroscopy in Solid Parahydrogen

Mario E. Fajardo

USAF Research Laboratory, AFRL/PRSP
Bldg. 8451, Edwards AFB, CA 93524-7680
mario.fajardo@edwards.af.mil

- * Cryosolid Propellants Team
- * HEDM Cryosolid Propellants Concept (Atoms in Solid Hydrogen)
- * Rapid Vapor Deposition of Transparent Parahydrogen (pH₂) Solids
- * B and Al Doped pH₂ Solids
- * High Res. IR Spectroscopy of Molecular Dopants in Solid pH₂
- * Summary

Cryosolid Propellants Team

Mario E. Fajardo, Michelle E. DeRose, and Simon Tam

- * Mario E. Fajardo, Michelle E. DeRose, and Simon Tam
Bill Larson (thermal B atom source)
- * Jeff Sheehy, Jerry Boatz, Peter Langhoff (in-house theory)
- * AFOSR Contractors:
 - P. Dagdigan @ Johns Hopkins: Al/H₂ & B/H₂ Complexes
 - M. Alexander @ U. Maryland: B/H₂ Interaction Potentials
 - G. Voth @ U. Utah: Path-Integral Monte Carlo Simulations
 - G. Scoles & K. Lehmann @ Princeton U.: Helium Clusters
- * External Collaborators:
 - T. Momose @ Kyoto U.: High Resolution IR Spectroscopy
- * Summer Visiting Professors:
 - R.J. Hinde @ U. Tennessee: Dopant-Induced IR Activity
 - D. Anderson @ U. Wyoming: Dopant IR Absorptions

Propellant Performance Figures of Merit

Specific Impulse, I_{sp} :

$I_{sp} \equiv$ (total impulse / propellant weight)

$= g_0 \langle v_{exh} \rangle$ “seconds”

$$\propto \sqrt{\frac{\langle \text{K.E.} \rangle}{m}} \propto \sqrt{\frac{\Delta H}{m}} = \sqrt{\Delta H_{sp}}$$

Density, ρ :

higher density \Rightarrow smaller & lighter tanks

\Rightarrow less aerodynamic drag

\Rightarrow condensed phase propellants

[G.P. Sutton, “Rocket Propulsion Elements” (Wiley, New York, 1992).]

“Revolutionary” vs. “Evolutionary”

HEDM Concepts


- * “Revolutionary” means better than LOX/LH₂:
LOX/LH₂ $\Delta H_{\text{sp}} = 12.6 \text{ MJ/kg}$ (3.0 kcal/g)
HEDM Target: $\Delta H_{\text{sp}} > 15.0 \text{ MJ/kg}$ (3.6 kcal/g)

- * Early (c1990) Revolutionary HEDM Concepts:
 - tetrahydrogen (H₄)
 - metastable triplet helium (He* and He₂*)
 - spin-polarized atomic hydrogen (H↑)
 - high-spin species (⁵CO)
 - dications (AB⁺⁺, ABC⁺⁺)

- ♥ “non-metallics” (e.g. O₄/H₂, N₄, N₈, N₂₀) N₅⁺!
- ♥ metallic hydrogen
- ♥ metal atoms and clusters in solid H₂

Cryosolid Propellants Concept

Use cryogenic solid hydrogen as a “packaging material” to store energetic species such as metal atoms and clusters.



Atom Additive Payoffs (5 % molar)

Sea level specific impulse, I_{sp} , in seconds (% change)

$$P_{\text{chamber}} = 1000 \text{ PSIA}, P_{\text{exhaust}} = 14.7 \text{ PSIA}$$

Additive	in standard state <u>M(5%)/LOX/H₂</u>	as atoms <u>M(5%)/LOX/H₂</u>	monoprop. <u>M(5%)/H₂</u>
none	403		
C	381 (-5%)	515 (+28%)	515 (+28%)
B	407 (+1%)	508 (+26%)	465 (+15%)
Be	427 (+6%)	493 (+22%)	
Si	400 (-1%)	460 (+14%)	
Al	407 (+1%)	454 (+13%)	
H	403	430 (+7%)	380 (-6%)
Li	404	428 (+6%)	
Mg	400 (-1%)	416 (+3%)	


Cryosolid Propellants Objectives

- * Make solid hydrogen samples (any size) containing 5% molar concentration of trapped energetic additives.
- * Measure absolute concentrations of energetic species.
- * Scale-up samples; produce $\sim 1 \text{ cm}^3$ samples in our lab.

Example: 5% Al/pH₂, $V = 1 \text{ cm}^3$
assume each Al atom replaces one H₂ molecule
 $\Rightarrow 58 \text{ mg Al} / 83 \text{ mg } ^1_0\text{H}_2$ (*see display item*)
 $\therefore \rho = 0.142 \text{ g/cm}^3$ (+100%)

Cryosolid Propellants Approach

- * Rapid vapor deposition of metal atom vapor and pre-cooled parahydrogen gas onto a liquid helium cooled substrate in vacuum.



Dopant Reactions within solid pH_2

* ideally:




isolated atoms



* in practice:



recombination



reaction



both



The Perils of Calorimetry

108

GEORGE C. PIMENTEL

TABLE IX
CONCENTRATIONS OF FREE RADICALS REPORTED

Radical	Matrix	Mole per cent radicals	Method of production and estimate ^a	Reference
O	O ₂	4-20	Gas, cal	Minkoff <i>et al.</i> (1959).
		<3	Gas, IR	Harvey and Bass (1958)
OH	Ca(OH) ₂	~1	Gas, cal	Broida and Lutes (1956)
N	N ₂	0.6	γ, ESR	R. Livingston ^b
		4	Gas, cal	Minkoff <i>et al.</i> (1959)
		0.2	Gas, cal	Broida and Lutes (1956)
		0.03	γ, ESR	Wall <i>et al.</i> (1959b)
		>0.03	Gas, cal	Fontana ^c (1958)
		0.01-0.04	Gas, MS	Matheson and Smaller (1955)
OH(?)	HCOOH	0.2	γ, ESR	Wall <i>et al.</i> (1959a)
CH ₃	CH ₄	0.14	γ, ESR	Wall <i>et al.</i> (1959a)
H	CH ₄	0.1	γ, ESR	Cole and Harding (1958)
N	NH ₃	0.1	Gas, ESR	Livingston <i>et al.</i> (1955)
H	HClO ₄ -H ₂ O	0.1	γ, ESR	Matheson and Smaller (1955)
H	H ₂ O	0.01	γ, ESR	Matheson and Smaller (1955)
H, NH ₂ (?)	NH ₃	0.01	γ, ESR	D. Ingram ^b
ROH	Alcohols	~0.01	UV, ESR	Wall <i>et al.</i> (1959a)
H	H ₂	0.0006	γ, ESR	


^a Abbreviations: gas = rapid condensation of gaseous radicals; γ = gamma ray *in situ* production; UV = photolytic *in situ* production; IR = infrared analysis; cal = calorimetry; MS = magnetic susceptibility.

^b Private communication.

^c Fontana, B. J. (1959). *J. Chem. Phys.* **31**, 148.


[A.M. Bass and H.P. Broida, "Formation and Trapping of Free Radicals" (Academic, New York, 1960).]

Experimental Diagram (c1993)



M.E. Fajardo, J. Chem. Phys. **98**, 110 (1993).

Transmission Spectrum of Li/nH_2 , $d \approx 10 \mu$



M.E. Fajardo, J. Chem. Phys. **98**, 110 (1993).

Optical Scattering in Solid Hydrogen

Crystal Growing and Quality (p. 81)

“There is a considerable art to growing hydrogen crystals of high quality. Good crystals are always grown slowly from the melt; a rapid freeze from the gas produces snow.”


Crystallite Light Scattering (p. 83)

“The reason that a good hydrogen crystal is so hard to see is its low refractive index...an estimated 1.16!

Yet a 1 mm-thick layer of hydrogen crystallites can be a completely opaque brown-black.”


[P.C. Souers, Hydrogen Properties for Fusion Energy (UC Press, Berkeley, 1986)]

ortho- and para-hydrogen




[L.F. Silveira, Rev. Mod. Phys. 52, 393 (1980)]

Rapid Vapor Deposition of Gram-Scale Optically Transparent pH_2 Solids (c1997)




M.E. Fajardo and S. Tam, J. Chem. Phys. **108**, 4237 (1998).
S. Tam and M.E. Fajardo, Rev. Sci. Instrum. **70**, 1926 (1999).

Electronic Spectroscopy of B/pH₂ ($d \approx 2$ nm)




S. Tam, M. MacIer, M.E. DeRose, and M.E. Fajardo, J. Chem. Phys. **113**, 9067 (2000).
[J.R. Krumrine, S. Jang, G.A. Voth, and M.H. Alexander, J. Chem. Phys. **113**, 9079 (2000)]

Photodynamics Cartoon for B/pH₂




S. Tam, M. MacIer, M.E. DeRose, and M.E. Fajardo, J. Chem. Phys. **113**, 9067 (2000).

Photobleaching of B/pH₂ Absorptions



S. Tam, M. Macler, M.E. DeRose, and M.E. Fajardo, J. Chem. Phys. **113**, 9067 (2000).

IR Absorption Spectra of B_2H_6/pH_2




S. Tam, M. MacIer, M.E. DeRose, and M.E. Fajardo, J. Chem. Phys. **113**, 9067 (2000).

Requirement for High Flux HEDM Sources

* A 5 % doping level, and a sample growth rate of 1 mm/hour, require a flux of $\Phi_{\text{HEDM}} \approx 3 \times 10^{16} \text{ \#/cm}^2\text{-s}$ at the deposition substrate. For Al atoms, this translates to a mass flux of 5.8 mg/cm²-hour, delivered to the deposition substrate.

* Began FY00 using miniature tungsten filament evaporation sources based on our FY99 effort to produce thermal B atoms.

* Total mass loadings of Al metal were $\sim 10 \text{ mg}$, just enough to detect trapped Al atoms in Ar; $\Phi_{\text{Al}} \sim 10^{11} \text{ \#/cm}^2\text{-s @ } R = 5 \text{ cm}$.



High Flux HEDM Sources

- * Purchased commercial Al evaporator; PBN crucible holds ≈ 10 g Al in horizontal orientation.
- * $T_{\max} = 1200\text{ }^{\circ}\text{C} \Rightarrow P_{\text{vap}}(\text{Al}) \approx 8 \times 10^{-3} \text{ torr} \Rightarrow \Phi_{\text{Al}} \approx 10^{18} \text{ \#/cm}^2\text{-s}$





Figure 1-3: Schematic of the EPI SUMO™ Effusion Cell.


UV Spectroscopy Al/pH₂

- * Al atom UV absorption saturates at high column densities.




Department of Chemistry, University of Wyoming, Laramie, WY, 20 April 2001

Recombination/reaction in Al/pH_2



Assignment of Al/pH₂ UV Absorptions



High Res. IR Spectroscopy in Solid pH_2





FIG. 1. Apparatus for the simultaneous spectroscopy of the infrared and Raman transitions. The nonlinearity of LiNbO_3 is used for the former and that of solid H_2 is used for the latter. D.M., dichroic mirror; S. A., spectrum analyzer; P. B., polarizer beamsplitter.

IR Absorption of 6 mm Thick pH_2 Solid

Non-observation of the $Q_1(0)$ transition demonstrates the absence of oH_2 impurities, and that the microscopic structure is not amorphous or porous.


Observation of $S_1(0)$ transition demonstrates the absence of inversion symmetry for some H_2 molecular environments.

[van Kranendonk and Gush, Phys. Lett. **1**, 22 (1962)]



M.E. Fajardo and S. Tam, J. Chem. Phys. **108**, 4237 (1998).

Raman Spectra of pH_2 Solids




Mixed hcp/fcc as-deposited structure, anneals to hcp; compare with: [G.W. Collins, et al., Phys. Rev. B **53**, 102 (1996)].

(d) sample in (c) warmed to 4.5 K.
(c) 4.5 mm sample as deposited at 3.3 K ($\Phi = 290$ mmol/hr).

(b) sample in (a) warmed to 4.5 K.
(a) 6 mm sample as deposited at 3.1 K ($\Phi = 200$ mmol/hr).


M.E. Fajardo and S. Tam,
J. Chem. Phys. **108**, 4237 (1998).

CH₄ Nuclear Spin Modifications




[M. Hepp, G. Winniewisser, and K.M.T. Yamada, J. Mol. Spectr. **164**, 311 (1994)]

ν_4 CH₄/pH₂ IR Absorptions (res = 0.01 cm⁻¹)




S. Tam, M.E. Fajardo, H. Katsuki, H. Hoshina, T. Wakabayashi, and T. Momose, J. Chem. Phys. **111**, 4191 (1999).

ν_4 CH₄/pH₂ IR Absorptions




- (a) Rapid Vapor Deposited sample: as-deposited at 2.4 K
- (b) Rapid Vapor Deposited sample: annealed to 4.8 K
- (c) Enclosed Cell Condensed sample: cooled to 4.8 K


v_4 CH_4/pH_2 Energy Levels



CH_4/pH_2 from Laser Ablation of Graphite




Co-operative IR absorptions



analysis in collaboration with D.T. Anderson, U. Wyoming and R.J. Hinde, U. Tennessee, Knoxville.

88 PPM HCl/pH₂



Gas Phase $(\text{HCl})_2$

High resolution, jet-cooled infrared spectroscopy of $(\text{HCl})_2$: Analysis of ν_1 and ν_2 HCl stretching fundamentals, interconversion tunneling, and mode-specific predissociation lifetimes

Michael D. Schuder,^{a)} Christopher M. Lovejoy,^{b)} Robert Lascola,^{c)} and David J. Nesbitt^{d)}
*Joint Institute for Laboratory Astrophysics, National Institute of Standards and Technology and
 University of Colorado, and the Department of Chemistry and Biochemistry, University of Colorado,
 Boulder, Colorado 80309*

(Received 5 April 1993; accepted 7 June 1993)





FIG. 1. Vibrationally averaged structure and internal coordinates for HCl dimer. The internuclear axis R connects the HCl centers of mass. The internal angles, θ_1 and θ_2 , are measured from the HCl centers of mass to the HCl bonds r . The torsion angle, $\phi = \phi_1 - \phi_2$, is shown at 180° (planar). The minimum energy configuration shown is for $\theta_1 = 16^\circ$, $\theta_2 = 87^\circ$ with $\phi_1 - \phi_2 = 180^\circ$. The HCl subunit on the left is referred to as the bonded HCl with an associated vibration labeled ν_2 . The proton on the other HCl is not involved with the hydrogen bond, and this subunit is referred to as the free HCl , with a vibration labeled ν_1 .

$(\text{HCl})_2/\text{pH}_2$ isotopomers



analysis in collaboration with D.T. Anderson, U. Wyoming.

HEDM Cryosolids Accomplishments

(a list of “things that’ll never work.”)

- * Trapped Li, B, Na, Mg, Al atoms in solid hydrogen at $T \approx 2$ K; attempts to demonstrate useful chemical energy storage still in progress
- * Demonstrated production of gram-scale optically transparent pH_2 solids by rapid vapor deposition.
- * Demonstrated that vapor deposited pH_2 solids are densest close-packed solids, NOT amorphous.
- * Demonstrated suitability of vapor deposited pH_2 solids as hosts for high resolution IR absorption spectroscopy of chemically interesting dopants; spectral assignments ongoing.
- * Generalized phenomena of dopant-induced and co-operative IR absorptions to chemically interesting dopants.